

## TONSILLECTOMY – ADULTS & CHILDREN

A watchful wait approach is more appropriate than tonsillectomy surgery for both children and adults with mild sore throats.

Fast track referral for specialist assessment and investigation for malignancy (which may include tonsillectomy for biopsy) or emergency presentations **are not restricted and are exclusions to this policy**. There are also rare indications, where as part of an underlying illness, it is good medical practice to perform a tonsillectomy.

### CRITERIA BASED ACCESS for the following indication only:

Recurrent peritonsillar abscess (quinsy), which has required a hospital admission or antibiotics/drainage

## PRIOR APPROVAL REQUIRED

### Prior Approval is required prior to referral for the following:

#### Tonsillitis

Before referral to secondary care, discuss with patient/parents or carers the benefits and risks of tonsillectomy vs. active monitoring. Referral for consideration of surgery will be funded if the following criteria are met:

The patient must have:

- Sore throats due to acute recurrent tonsillitis where the episodes are disabling and cause significant functional impairment.

#### WITH

- Seven or more well documented clinically significant, adequately treated sore throats in one year.

#### OR

- Five or more such episodes in each of the preceding 2 years

#### OR

- Three or more such episodes in each of the preceding 3 years

**Clinical details (e.g., dates when antibiotics were prescribed) are to be included in the prior approval submission.**

Reference:	Policy Name	Review Date	Version
BSW-CP005	Tonsillectomy	November 2024	V1.3 November 2021

## PRIOR APPROVAL REQUIRED

**Obstructive sleep apnoea / Obstructive sleep disordered breathing in children (aged 18 and under)**

Adenotonsillectomy may be considered as part of treatment for obstructive sleep disordered breathing in children.

Obstructive Sleep Apnoea in children should be diagnosed clinically:

- A history of witnessed sleep apnoeic attacks in the presence of snoring.

**AND**

- The presence of large and obstructive tonsils

**AND**

- Failure to thrive (documented impact on development, behaviour and quality of life, height and weight, hyperactivity, daytime somnolence)

**AND**

- Absence of any neurological deficit which might cause Central Sleep Apnoea.

(A sleep study i.e., polysomnography is not required for the diagnosis of Obstructive Sleep Apnoea in children.)

**Of note** - children with simple snoring without symptoms or signs of apnoea are unlikely to benefit from adenotonsillectomy.

**Obstructive sleep apnoea/hypopnoea syndrome in adults**

NICE guidance recommends that tonsillectomy may be considered for people with OSAHS who:

- Have large obstructive tonsils

**AND**

- A body mass index (BMI) of less than 35 kg/m<sup>2</sup>.

## INTERVENTIONS NOT NORMALLY FUNDED

### Exceptional Funding Required

- Tonsillectomy procedures for tonsil stones or halitosis.
- Tonsillectomy for simple snoring (without symptoms or signs of apnoea)
- Asymmetry of the tonsils where malignancy is not suspected.

Reference:	Policy Name	Review Date	Version
BSW-CP005	Tonsillectomy	November 2024	V1.3 November 2021